

The Synoptic Swift Synergy – Catching Gamma-Ray Bursts Before They Fly

Jeremy S. Heyl^{1,2}

ABSTRACT

The advent of large panoramic photometric surveys of the sky offers the possibility of exploring the association of gamma-ray bursts (GRBs) with supernovae. To date, a few gamma-ray bursts have been connected possibly with supernovae: GRB 980425 – SN 1998bw, GRB 011121 – SN 2001ke, GRB 970228 and GRB 980326. A combination of a large detection rate of GRBs and rapid coverage of a large portion of the sky to faint magnitude limits offers the possibility of detecting a supernova preceding an associated GRB or at least placing limits on the rate of association between these two phenomena and the time delay between them. This would provide important constraints on theoretical models for gamma-ray bursts.

Subject headings: gamma rays: bursts

1. Introduction

The gamma-ray burst GRB 021004 was detected by HETE II at 12:06 UT on the 4th of October 2002 (Shirasaki et al. 2002). Observations after about 9 minutes from the trigger revealed a fading optical transient (Fox 2002), which was densely sampled in several bands, especially at early times. The afterglow of GRB 021004 has shown several unusual features (Mirabal et al. 2002; Salamanca et al. 2002; Moller et al. 2002; Bersier et al. 2003). Perhaps its most unique feature was that the field had been observed shortly before the gamma-ray burst itself was detected. Wood-Vasey et al. (2002) give a limiting unfiltered magnitude of 21.4 for observations on the day before the burst and 22.3 integrated over the year before the burst.

Astronomy is on the threshold of a new era where large portions of the sky are surveyed deeply and regularly. The question arises what is likelihood of getting photometry of a

¹Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138; jhey1@cfa.harvard.edu

²Chandra Fellow

gamma-ray burst precursor, specifically if supernovae precede gamma-ray bursts as in the supranova model (Vietri & Stella 1998)? Although the flux upper limits for GRB 021004 are not stringent enough to constrain theoretical models of gamma-ray bursts, the high-burst localization rate of Swift combined with the fast sky coverage of the Sloan Digital Sky Survey (SDSS) and later Pan-STARRS, LSST and the Supernova-Acceleration Probe (SNAP) could provide important constraints on gamma-ray burst precursors.

During the first and second years of operation of the Swift mission, SDSS will scan approximately 3000 square degrees (or 7% of the sky) each year (SDSS Collaboration 2001). Over this area it will detect point sources down to $R \approx 23.2$. If Swift or subsequent missions are operational in 2006, the Pan-STARRS program will observe 20,000 square degrees every four days (or 50% of the sky) to a limiting magnitude of $R \approx 24.2$ (Kaiser et al. 2002). Finally, potentially beginning in 2010, SNAP will cover 15 square degrees every four days with each observation reaching a limiting magnitude of $R \approx 28$ (Kim et al. 2002), coadding observations over a month would go one magnitude deeper. The SNAP lensing survey will cover 300 square degrees over five months to a similar limiting magnitude.

Long gamma-ray bursts are thought to be associated with the collapse of a massive star, a supernova. Specifically, in the collapsar model, the formation of a black hole in the center of the star results in relativistic jets which pierce the envelope of the star (MacFadyen & Woosley 1999). Along the axis of the jets, the collapsing star appears as a gamma-ray burst, and the supernova reaches its peak a few weeks after the GRB. Vietri & Stella (1998) proposed an alternative model in which the gamma-ray burst accompanies the delayed collapse of a quickly spinning neutron star which is more massive than the maximum mass of a non-rotating neutron star. The neutron star may take several months or years after the supernova to spin down to the critical frequency and collapse.

In this *Letter*, I will estimate the number of gamma-ray burst events with photometry which overlaps on the sky but shortly precedes in epoch from SDSS and other surveys and compare the flux limits with the expected flux from a supernova which may precede the gamma-ray burst.

2. Gamma-Ray Burst Overlap with Future Surveys

To calculate how often sufficiently deep photometry will precede the observation of a gamma-ray burst on the sky, several ingredients are required: a model for the spectral-energy distribution as a function of time of a supernova associated with a GRB, an estimate of the luminosity-rate function of GRBs as a function of redshift ($\dot{\phi}(z, L)$), a model for the field of

view of the gamma-ray burst detector ($\Omega_{\text{GRB}} = 2$ for Swift) and its detection threshold (P_1) and the rate of sky coverage of the photometric program ($\mathcal{R}_{\text{photo}}$) and its detection threshold (R_{lim}). Porciani & Madau (2001) provide models for $\dot{\phi}(z, L) \equiv R_{\text{GRB}}(z)\psi(L)$. The rate of GRBs, $R_{\text{GRB}}(z)$, is taken to be proportional to the star-formation rate, and the luminosity function of GRBs, $\psi(L)$, is constrained by the BATSE GRB number counts. The rate of overlapping photometry is given by the product of the rate of sky coverage with a integral over the assumed cosmological distribution of GRBs,

$$\frac{dN_{\text{overlap}}}{dt}(P > P_1, R < R_{\text{lim}}) = \frac{\mathcal{R}_{\text{photo}}\Delta t}{4\pi} \frac{dN_{\text{total}}}{dt}(P > P_1, R < R_{\text{lim}}) \quad (1)$$

$$\frac{dN_{\text{total}}}{dt}(P > P_1, R < R_{\text{lim}}) = \frac{\Omega_{\text{GRB}}}{4\pi} \int_0^{z: R(z)=R_{\text{lim}}} dz \int_{L(P_1, z)}^{\infty} dL \frac{dV(z)}{dz} \frac{\dot{\phi}(z, L)}{1+z}. \quad (2)$$

For lack of a better model for the evolution of a supernova associated with a GRB, I will assume that SN2001ke (Garnavich et al. 2003) is a prototype for this class, and furthermore that a supernovae associated with a GRB maintains its peak brightness for a period $\Delta t = 14(1+z)$ days in the observer’s frame and otherwise it is undetectable (see Reichart 1999; Bloom et al. 1999, for other GRB-associated supernovae). It is reasonable to use the median value of z for GRBs whose associated supernova are brighter than the magnitude limit of the particular photometric survey. However, to be highly conservative, I will take $\Delta t = 14$ days to calculate the rate of overlap.

If a survey covers the same area of sky more often than once per interval Δt such as the SNAP supernovae search and Pan-STARRS, the rate of sky coverage $\mathcal{R}_{\text{photo}}$ should only account for the first visit in each period Δt ; for example $\mathcal{R}_{\text{photo}}$ for Pan-STARRS is 2π per fourteen days. The additional visits during each fortnight do not increase $\mathcal{R}_{\text{photo}}$ but they do allow the survey to probe deeper by coadding the successive images.

According to the original supranova model (Vietri & Stella 1998), the supernova may reach its peak at any time up to several years before the GRB, so this calculation implicitly assumes that both the GRB survey and the photometric survey will be operating at the appropriate times. $L(P_1, z)$ is the luminosity of a GRB at a redshift z which is detected at a count-rate of P_1 , and $R(z)$ is the R -band apparent magnitude of a GRB-associated supernovae at a redshift z . Both of these functions include the k -correction (Hogg 1999) and assume the cosmographic parameters, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$. Fig. 1 plots the number of GRBs detected by Swift per year whose associated supernova would be brighter at its peak than a particular R -band magnitude.

The results shown in Fig. 1 assume the SF1 model of Porciani & Madau (2001). This model provides a conservative lower limit for the overlap. It predicts that Swift will localize about 110 bursts per year — the more generous estimates range up to 300 bursts per year




Fig. 1.— The number of GRB-associated supernova brighter than a given R -magnitude. The lines show the cumulative contribution of GRBs above given flux limits. The right panel shows the entire distribution while the left panel focusses on the bright end. From bottom to top, only the supernovae associated with GRBs whose peak flux is above $10^{0.9}$, $10^{0.6}$, $10^{0.5}$, $10^{0.4}$, $10^{0.3}$, $10^{0.1}$, 1 , $10^{0.2}$, $10^{-0.25}$, $10^{-0.45}$, $10^{-0.5}$, $10^{-0.7}$ and $10^{-0.75}$ photons per square centimeter per second. See Porciani & Madau (2001) for further details.

(Myers 2002). Furthermore, this model predicts that the bursts detected will be at higher redshifts than other models, so the accompanying supernovae will be fainter and more difficult to detect.

Table 1 gives the overlap rate between various photometric surveys and the Swift GRB localization mission. What is striking is that the shallow but wide Pan-STARRS and LSST surveys will perform much better than any of the other surveys. Furthermore, if supernovae precede GRBs, Pan-STARRS and LSST each will detect nearly ten GRB-associated supernovae per year. If it finds none, it would place severe constraints on the supernova model for GRBs. It must be emphasized that this rate of overlap is extremely conservative. It assumes a low Swift burst localization rate and a distribution of GRBs skewed to high redshift (therefore, faint associated supernovae). The actual rate of overlap will probably be higher if both programs operate simultaneously. Furthermore, Swift will generate a catalog of burst positions and redshifts. One will be able to cross-correlate *a posteriori* this catalog with earlier Pan-STARRS or LSST observations and exclude the appearance of transients to $R \approx 25$ over a wide range of epochs preceding the burst yielding definitive constraints on GRB-progenitors independent of assumptions about the GRB luminosity function and its evolution.

This calculation of the overlap rate assumes that either the GRB localization program or the photometric survey studies random portions of the sky. In fact both the Swift mission and all of the photometric surveys avoid studying the region of the sky near the sun. Although the average rate of overlap over a year is given by the formulae above and the values in the tables, the chance of detecting the supernova associated with GRB is somewhat higher than average if the supernova precedes the GRB by less than three months or between nine and fifteen months. If the supernova precedes the GRB by six to nine months, the chance of detecting it is somewhat lower than average. However, this seasonal variation is smaller than the uncertainties in the GRB luminosity function.

3. Discussion

The philosophy employed for finding gamma-ray burst precursors is somewhat different than what is necessary for finding supernovae or microlensing events. Because the precursors will be sought after the gamma-ray burst is detected and localized, it is not necessary to have more than one epoch of data from the particular region of sky before the burst. Even a single epoch would yield important constraints. Furthermore, unless the cadence of the observations is sufficiently low (no more than biweekly), the repeated observations of the same patch of sky do not improve the chances of catching a precursor (unless one coadds

the data to probe deeper), because supernova typically evolve over the course of weeks. Consequently, although supernova and microlensing surveys have a large data rate of high quality photometry, because of their relative lack of sky coverage and depth they do not contribute much to the detection rate of precursors. From another point of view, only a small fraction ($< 10^{-4}$) of supernovae result in GRBs directed toward us, so one would typically have to find at least $10^4(4\pi/\Omega_{\text{GRB}})$ supernovae in a blind search before finding a single GRB-associated supernova.

The best bets are the large deep wide surveys of the sky. SDSS is the prototype and Pan-STARRS and LSST should deliver results. There is a small possibility that SDSS will catch a supernova before a gamma-ray burst providing important evidence for the supernova model for gamma-ray bursts (it may have done so already). Pan-STARRS or LSST if it overlaps with a high-localization-rate GRB mission such as Swift will be able to provide important constraints on gamma-ray-burst models. Specifically, it will be able to exclude the possibility that GRBs follow supernovae within a year.

I would like to acknowledge useful discussions with Kris Stanek, Robert Lupton and Bob Kirshner and helpful suggestions from the anonymous referee. I was supported by the Chandra Postdoctoral Fellowship Award # PF0-10015 issued by the Chandra X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-39073.

REFERENCES

- Bersier, D. et al. 2003, ApJ, 584, L43
- Bloom, J. S. et al. 1999, Nature, 401, 453
- Fox, D. 2002, GCN, 1564
- Garnavich, P. M. et al. 2003, ApJ, 582, 924
- Hogg, D. 1999, astro-ph/9905116
- Kaiser, N., Boesgaard, H., Heasley, J., Hodapp, K., Jewitt, D., Kudritzki, R.-P., Kuhn, J., Luppino, G., Magnier, E., Onaka, P., Pickles, A., Simon, T., Szapudi, I., Tholen, D., & Tonry, J. 2002, POI: The Panoramic Optical Imager, Tech. rep., Institute for Astronomy

- Kim, A. et al. 2002, Wide-Field Surveys with the SNAP Mission, Tech. rep., Lawrence Berkeley Laboratory
- MacFadyen, A. I. & Woosley, S. E. 1999, *ApJ*, 524, 262
- Mirabal, N., Armstrong, E. K., Halpern, J. P., & Kemp, J. 2002, GCN, 1602
- Moller, P. et al. 2002, *A&A*, in press (astro-ph/0210654)
- Myers, J. D. 2002, Swift: Catching Gamma-Ray Bursts on the Fly, Tech. rep., NASA, <http://swift.nasa.gov>
- Porciani, C. & Madau, P. 2001, *ApJ*, 548, 522
- Reichart, D. E. 1999, *ApJ*, 521, L111
- Salamanca, I., Rol, E., Wijers, R., Ellison, S., Kaper, L., & Tanvir, N. 2002, GCN, 1611
- SDSS Collaboration. 2001, Five-Year Baseline Plan for SDSS Operations, Tech. rep., Sloan Digital Sky Survey
- Shirasaki, Y. et al. 2002, GCN, 1565
- Vietri, M. & Stella, L. 1998, *ApJ*, 507, L45
- Wood-Vasey, W. M., Aldering, G., Lee, B. C., Helin, E. F., Pravdo, S., Hicks, M., & Lawrence, K. 2002, GCN, 1572

Table 1. Present and Future Large-Scale Photometric Surveys

Survey	R_{lim}	z_{max}	z_{med}	$\mathcal{R}_{\text{photo}}\Delta t$	$dN_{\text{total}}/dt[\text{y}^{-1}]$	$dN_{\text{overlap}}/dt[\text{y}^{-1}]$
SDSS	23.2	0.56	0.48	0.035	1.9	0.0052
Pan-STARSS (single)	24.2	0.78	0.66	6.3	7.1	3.6
Pan-STARSS (coadded)	25.0	1.00	0.83	6.3	17.	8.8
LSST (single)	24.5	0.86	0.72	6.3	10.	5.1
LSST (coadded)	25.1	1.04	0.85	6.3	19.	9.8
SNAP SN (single)	28.0	2.59	1.45	0.0046	98.	0.036
SNAP SN (coadded)	28.8	3.35	1.50	0.0046	110.	0.039
SNAP lensing	28.0	2.59	1.45	0.0091	98.	0.071